

ON COBWEB POSETS AND THEIR COMBINATORIALLY ADMISSIBLE
SEQUENCES

Andrzej Krzysztof Kwaśniewski

Member of the Institute of Combinatorics and its Applications
High School of Mathematics and Applied Informatics
Kamienna 17, PL-15-021 Białystok, Poland
e-mail: kwandr@gmail.com

SUMMARY

The purpose of this article is to pose three computational problems which are quite easily formulated for the new class of directed acyclic graphs interpreted as Hasse diagrams. The problems posed are not yet solved though are of crucial importance for the vast class of new partially ordered sets with joint combinatorial interpretation. These so called cobweb posets - are relatives of Fibonacci tree and are labeled by specific number sequences - natural numbers sequence and Fibonacci sequence included. One presents here also a join combinatorial interpretation of those posets' F -nomial coefficients which are computed with the so called cobweb admissible sequences. Cobweb posets and their natural subposets are graded posets, sometimes called a ranked posets. They are vertex partitioned into such antichains Φ_n (where n is a nonnegative integer) that for each Φ_n , all of the elements covering x are in Φ_{n+1} and all the elements covered by x are in Φ_n . We shall call the Φ_n the n -th level. The cobweb posets may be identified with a chain of di-bicliques i.e. by definition - a chain of complete bipartite one direction digraphs. Any chain of relations is therefore obtainable from the cobweb poset chain of complete relations via deleting arcs in di-bicliques of the complete relations chain.

Key Words: acyclic digraphs, tilings, special number sequences, binomial-like coefficients.

AMS Classification Numbers: 06A07 ,05C70,05C75, 11B39.

affiliated to The Internet Gian-Carlo Polish Seminar:

http://ii.uwb.edu.pl/akk/sem/sem_rota.htm

Published in : Adv. Studies Contemp. Math. Vol. **18** No 1, (2009), 17-32.

1 Introduction

In computer science a directed acyclic graph, also called DAG, is a directed graph with no directed cycles. Utility of DAG's is widely known and appreciated. Some algorithms - for example, search algorithms - become simpler when applied to DAGs. DAGs considered as a generalization of trees have a lot of applications in computer science, bioinformatics, physics and many natural activities of humanity and nature. For example in information categorization systems, such as folders in a computer or in Serializability Theory of Transaction Processing Systems and many others. Here we introduce specific DAGs as generalization of trees being inspired by algorithm of the Fibonacci tree growth. For

any given natural numbers valued sequence the graded (layered) cobweb posets' DAGs are equivalently representations of a chain of binary relations. Every relation of the cobweb poset chain is biunivocally represented by the uniquely designated **complete** bipartite digraph-a digraph which is a di-biclique designated by the very given sequence. The cobweb poset is then to be identified with a chain of di-bicliques i.e. by definition - a chain of complete bipartite one direction digraphs. Any chain of relations is therefore obtainable from the cobweb poset chain of complete relations via deleting arcs (arrows) in di-bicliques. Let us underline it again : *any chain of relations is obtainable from the cobweb poset chain of complete relations via deleting arcs in di-bicliques of the complete relations chain.* For that to see note that any relation R_k as a subset of $A_k \times A_{k+1}$ is represented by a one-direction bipartite digraph D_k . A "complete relation" C_k by definition is identified with its one direction di-biclique graph $d - B_k$. Any R_k is a subset of C_k . Correspondingly one direction digraph D_k is a subgraph of an one direction digraph of $d - B_k$.

The one direction digraph of $d - B_k$ is called since now on **the di-biclique** i.e. by definition - a complete bipartite one direction digraph. Another words: cobweb poset defining di-bicliques are links of a complete relations' chain.

Because of that the cobweb posets in the family of all chains of relations unavoidably is of principle importance being the most overwhelming case of relations' infinite chains or their finite parts (i.e. vide - subposets).

The intuitively transparent names used above (a chain of di-bicliques etc.) are to be the names of defined objects in what follows. These are natural correspondents to their undirected graphs relatives as we can view a directed graph as an undirected graph with arrowheads added.

The purpose of this note is to put several questions intriguing on their own apart from being fundamental for a new class of DAGs introduced below. Specifically this concerns problems which arise naturally in connection with a new join combinatorial interpretation of all classical $F - nomial$ coefficients - Newton binomial, Gaussian q -binomial and Fibonomial coefficients included. This note is based on [8,9] from which definitions and description of these new DAG's are quoted for the sake of self consistency. Applications of new cobweb posets' originated Whitney numbers from [9] such as extended Stirling or Bell numbers are expected to be of at least such a significance in applications to linear algebra of formal series as Stirling numbers, Bell numbers or their q -extended correspondent already are in the so called coherent states physics (see [13] for abundant references on the subject). Quantum coherent states physics is of course a linear theory with its principle of states' superposition.

The problem to be the next. As cobweb subposets P_n are vertex partitioned into antichains Φ_r for $r = 0, 1, \dots, n$ which we call levels - a question of canonical importance arises. Let $\{P_n\}_{n \geq 0}$ be the sequence of finite cobweb subposets (see below). What is the form and properties of $\{P_n\}_{n \geq 0}$'s characteristic polynomials $\{\rho_n(\lambda)\}_{n \geq 0}$ [15,17]? For example - are these related to umbra polynomials? What are recurrence relations defining the $\{\rho_n(\lambda)\}_{n \geq 0}$ family ?

1.1. Partially Ordered Sets - Elementary Information.

Let us recall indispensable notions thus establishing notation and terminology.

Definition 1 . A partially ordered set (poset) is an ordered pair $\langle P, \leq \rangle$, where

P is a set and $\leq$ is a partial order on P . Naturally $x < y \Leftrightarrow x \leq y \wedge x \neq y$.

Definition 2 . An element z covers an element x provided that there exists no third element y in between i.e. such y that $x < y < z$. In this case z is called an upper cover of x .

A graphical rendering of a partially ordered set (poset) is being displayed via the cover relation of the partially ordered set (poset) with an implied upward orientation by convention. It represents the Hasse Diagram of the poset [3].

Definition 3 . Hasse diagram is a graph constructed as follows. A point is drawn for each element of the set P and line segments are drawn between these points according to the two rules: **1.** If $x \leq y$ then the point corresponding to x appears lower in the drawing than the point corresponding to y . **2.** The edge between the points corresponding to elements x and y is included in the graph if and only if x covers y or y covers x .

In order to make the reading interactive one recommends to go through preliminary guiding exercises (cf. [3,2]).

Exercise 1. Draw the Hasse diagrams for Boolean algebras of orders $n = 2, 3, 4$, and 5 .

Exercise 2. Invent Hasse diagram for Fibonacci poset $\langle P, \leq \rangle$. Its subsequent generations Φ_s of elements $x, y \in P$ are designated by Fibonacci sequence as $x < y$ iff if x is an ancestor of y . $F_0 = 0$ corresponds to the empty root $\{\emptyset\}$.

Example. Gallery of Posets by Curtis Greene may be found in [2]. Compare therein the bottom levels of a Young-Fibonacci lattice, introduced by Richard Stanley with cobweb posets defined here below.

1.2. Computation and Characterizing Problems.

In the next section we define cobweb posets and their examples are given [8,9]. A join combinatorial interpretation of cobweb posets' characteristic binomial-like coefficients is provided too. This simultaneously means join combinatorial interpretation of fibonomial coefficients and all incidence coefficients of reduced incidence algebras of full binomial type [16].

Finally we formulate three problems (to be explained on the way): characterization and/or computation of cobweb admissible sequences **Problem 1**, cobweb layers partition characterization and/or computation **Problem 2** and the GCD-morphic sequences characterizations and/or computation **Problem 3** - all three interesting on their own.

2 Cobweb posets - presentation, examples and combinatorial interpretation

2.1. F - nomial coefficients.

Given any sequence $\{F_n\}_{n \geq 0}$ of nonzero reals ($F_0 = 0$ being sometimes acceptable as $0! = F_0! = 1$.) one defines its corresponding binomial-like F - nomial coefficients as in Ward's Calculus of sequences [18] as follows.

Definition 4 . $(n_F \equiv F_n \neq 0, \quad n > 0)$

$$\binom{n}{k}_F = \frac{F_n!}{F_k! F_{n-k}!} \equiv \frac{n_F^k}{k_F!}, \quad n_F! \equiv n_F(n-1)_F(n-2)_F(n-3)_F \dots 2_F 1_F;$$

$$0_F! = 1; \quad n_F^k = n_F(n-1)_F \dots (n-k+1)_F.$$

We have made above an analogy driven identifications in the spirit of Ward's Calculus of sequences [18]. Identification $n_F \equiv F_n$ is the notation used in extended Fibonomial Calculus case [10,11,12,6] being also there inspiring as n_F mimics n_q established notation for Gaussian integers exploited in much elaborated family of various applications including quantum physics (see [10,9,13] and references therein).

2.2. Cobweb infinite posets. Cobweb infinite posets Π are designated uniquely by any sequence of natural numbers $F = \{n_F\}_{n \geq 0}$ with the one root convention ($F_0 = 1$) and are by construction endowed with a kind of self-similarity property. (In Fibonacci case $F_0 = 0$ corresponds to the empty root $\{\emptyset\}$). Given any such sequence $\{F_n\}_{n \geq 0} \equiv \{n_F\}_{n \geq 0}$ of positive integers we define the partially ordered, graded infinite poset Π - called a cobweb poset - as follows. Its vertices are labeled by pairs of coordinates: $\langle i, j \rangle \in N \times N_0$ where N stays for natural numbers while N_0 denotes the nonnegative integers. Vertices show up in layers of $N \times N_0$ grid along the recurrently emerging subsequent s -th levels Φ_s ("generations") where $s \in N_0$.

Definition 5 . $\Phi_s = \{\langle j, s \rangle \mid 1 \leq j \leq s_F\}, s \in N_0$.

Definition 6 .

$$\Pi = \langle P, E \rangle, \quad P = \bigcup_{0 \leq p} \Phi_p, \quad E = \{\langle \langle j, p \rangle, \langle q, (p+1) \rangle \rangle\} \bigcup \{\langle \langle 1, 0 \rangle, \langle 1, 1 \rangle \rangle\},$$

$$1 \leq j \leq p_F, 1 \leq q \leq (p+1)_F.$$

We shall refer to Φ_s as to the set of vertices at the s -th level. The population of the k -th level ("generation") counts k_F different member vertices for $k > 0$ and one for $k = 0$. Here down a disposal of vertices on Φ_k levels is visualized for the

case of Fibonacci sequence. $F_0 = 0$ corresponds to the empty root $\{\emptyset\}$. On the account of the above definition we shall sometimes irrelevantly represent $\Pi = \langle P, E \rangle$, just by its vertex set P partitioned into countable family of antichains (levels) uniquely designated by the F -sequence choice. The same facility is to be used for subposets on the grounds of a legal abuse of perfectionism.

-----and-----so-----on-----up-----↑-----
 *****10-th-level
 *****9-th-level
 *****8-th-level
 *****7-th-level
 *****6-th-level
 *****5-th-level
 *****4-th-level
 *****3-rd-level

$\star$ ----- 2-nd-level
 $\star$ ----- 1-st-level
 $\emptyset$ ----- 0-th-level

Figure 0. The s - th levels in $N \times N_0$

Accompanying to the set P of vertices the set E of edges - we obtain the Hasse diagram. We may picture the partially ordered infinite set P from the Definition 6 with help of the sub-poset set P_m (rooted at F_0 level of the poset) to be continued then ad infinitum in now obvious way as seen from the figures Fig.1 – Fig.5 of P_m cobweb posets below. These non-trees look like Fibonacci-type trees with a specific “cobweb”.




Fig.1. Display of Natural numbers' cobweb poset.




Fig.2. Display of Even Natural numbers' cobweb poset.




Fig3. Display of Odd natural numbers' cobweb poset.




Fig.4. Display of divisible by 3 natural numbers' cobweb poset.




Fig.5. Display of Fibonacci numbers' cobweb poset.

Compare this with the bottom levels of a Young-*Fibonacci lattice*, introduced by Richard Stanley- in Curtis Greene's gallery of posets [2].

As seen above - for example the Fig.5. displays the rule of the construction of the Fibonacci cobweb poset. It is being visualized clearly while defining this *non-lattice* cobweb poset P with help of its incidence matrix ζ [16]. The incidence

matrix ζ representing uniquely just this cobweb poset P exhibits (see below) a staircase structure characteristic to Hasse diagrams of all cobweb posets.

$$\begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \dots \end{bmatrix}$$

Figure 6. The incidence matrix ζ for the Fibonacci cobweb poset

Note that the knowledge of ζ matrix explicit form enables one to count via standard algorithms [16] the Möbius matrix $\mu = \zeta^{-1}$ and other typical elements of incidence algebra perfectly suitable for calculating number of chains, of maximal chains etc. in finite sub-posets of P . (All elements of the corresponding real incidence algebra are then given by a matrix of the Fig.6 with 1's replaced by arbitrary real numbers).

Cobweb posets as complete bipartite digraph sequences.

As announced earlier the cobweb posets and their natural subposets P_n are graded posets. They are vertex partitioned into antichains Φ_k for $k = 0, 1, \dots, r, \dots$ (where r is a nonnegative integer) such that for each Φ_r , all of the elements covering x are in Φ_{r+1} and all the elements covered by x are in Φ_r . We shall call the Φ_n the n -th-level. P_n is then $(n+1)$ level ranked poset. We are now in a position to observe that the cobweb posets may be identified with a chain of di-bicliques i.e. by definition - a chain of complete bipartite one direction digraphs. This is outstanding property as then any chain of relations is obtainable from the corresponding cobweb poset chain of complete relations just by deleting arcs in di-bicliques of this complete relations chain. Indeed. For any given natural numbers valued sequence the graded (layered) cobweb posets' DAGs are equivalently representations of a chain of binary relations. Every relation of the cobweb poset chain is bi-univocally represented by the uniquely designated **complete** bipartite digraph - a digraph which is a di-biclique designated by the very given sequence. The cobweb poset may be therefore identified with a chain of di-bicliques i.e. by definition - a chain of complete bipartite one direction digraphs. Say it again, any chain of relations is obtainable from the cobweb poset chain of complete relations via deleting arcs (arrows) in chains' di-bicliques elements.

The above intuitively transparent names (a chain of di-cliques etc.) are the names of the below defined objects. These objects are natural correspondents of their undirected graphs relatives as we can view a directed graph as an undirected graph with arrowheads added. Let us start with primary notions remembering that any cobweb subposet P_k is a DAG of course.

A bipartite digraph is a digraph whose vertices can be divided into two disjoint sets V_1 and V_2 such that every arc connects a vertex in V_1 and a vertex in V_2 . Note that there is no arc between two vertices in the same independent set V_1 or V_2 . No two nodes of the same partition set are adjacent.

A one direction bipartite digraph is a bipartite digraph such that every arc originates at a node in V_1 and terminates at a node in V_2 . The extension of "being one direction" to k -partite digraphs is automatic. Note that there is no arc between two vertices in the same set. Intuitively - one may color the nodes of a bipartite digraph black and blue such that no arc exists between like colors. A k -partite not necessarily one direction digraph D is obtained from a k -partite undirected k -partite graph G by replacing every edge xy of G with the arc $\langle xy \rangle$, arc $\langle yx \rangle$ or both $\langle xy \rangle$ and $\langle yx \rangle$. The partite sets of D are the partite sets of G .

Definition 7 . A simple directed graph $G = (V, E)$ is called bipartite if there exists a partition $V = V_1 + V_2$ of the vertex set V so that every edge (arc) in E is incident with v_1 and v_2 for some v_1 in V_1 and v_2 in V_2 . It is complete if any node from V_1 is adjacent to all nodes of V_2 .

We shall denote our special case bipartite one direction digraphs as follows $G = (V_1 + V_2, E) \equiv L_{F_k, F_{k+1}}$ to inform that the partition has parts V_1 and V_2 where $|V_1| = F_k$ and $|V_2| = F_{k+1}$.

Notation. The special k - partite $\equiv k$ - level one direction digraphs considered here shall be denoted by the corresponding symbol $L_{p,q,\dots,r}$. The complete k -partite $\equiv k$ -level one direction complete digraph is coded $K_{p,q,\dots,r}$ as in the non directed case (with no place for confusion because only one direction digraphs are to be considered in what follows). $L_{F_k, F_{k+1}, \dots, F_n} \equiv \langle \Phi_k \rightarrow \Phi_n \rangle$ denotes $(n - k + 1)$ - partite one direction digraph $\equiv (n - k + 1)$ - level one direction digraph $n \geq k$ whose partition has (antichains from Π) the parts $\Phi_k, \Phi_{k+1}, \dots, \Phi_n$, $|\Phi_k| = F_k, |\Phi_{k+1}| = F_{k+1}, |\Phi_n| = F_n$. Any cobweb one-layer $\langle \Phi_k \rightarrow \Phi_{k+1} \rangle$, $k < n$, $k, n \in N \cup \{0\} \equiv Z_{\geq}$ is a complete bipartite one direction digraph $L_{F_k, F_{k+1}} = K_{F_k, F_{k+1}}$ i.e. by definition it is the *di-biclique*. The cobweb one-layer vertices constitute bipartite set $V(\langle \Phi_k \rightarrow \Phi_{k+1} \rangle) = \Phi_k + \Phi_{k+1}$ while edges are all those arcs incident with the two antichains nodes in the poset Π graph representation.

Any cobweb subposet $P_n \equiv \langle \Phi_0 \rightarrow \Phi_n \rangle \equiv L_{F_0, F_1, \dots, F_n}$, $n \geq 0$ is a $(n + 1)$ -partite $= (n + 1)$ -level one direction digraph. We shall keep on calling a complete bipartite one direction digraph a **di-biclique** because it is a special kind of bipartite one direction digraph, whose every vertex of the first set is connected by an arc originated in this very node to **every** vertex of the second set of the given bi-partition. Any cobweb layer $\langle \Phi_k \rightarrow \Phi_n \rangle$, $k \leq n$, $k, n \in N \cup \{0\} \equiv Z_{\geq}$ is a one direction the $(n - k + 1)$ -partite $\equiv (n - k + 1)$ -level one direction DAG $L_{F_k, F_{k+1}, \dots, F_n}$ with an additional defining property: it is a chain of di-bicliques for $k < n$. Since now on we shall identify both:

$$\langle \Phi_k \rightarrow \Phi_n \rangle = L_{F_k, F_{k+1}, \dots, F_n}.$$

Note: $P_n \equiv \langle \Phi_0 \rightarrow \Phi_n \rangle = L_{F_0, F_{k+1}, \dots, F_n}$, $n \geq 0$ and $\langle \Phi_k \rightarrow \Phi_n \rangle^\neg = K_{F_k, F_{k+1}, \dots, F_n}$ for $n > k + 1$.

Observation.

$$|E(\langle \Phi_k \rightarrow \Phi_{k+m} \rangle)| = \sum_{i=0}^{m-1} F_{k+i} F_{k+i+1},$$

where $E(G)$ denotes the set of edges of a graph G (arcs of a digraph G).

The following property (*)

$$(*) \quad \langle \Phi_k \rightarrow \Phi_{k+1} \rangle \equiv L_{F_k, F_{k+1}} = K_{F_k, F_{k+1}}, \quad k = 0, 1, 2, \dots$$

i.e. $L_{F_k, F_{k+1}}$ is a di-biclique for $k = 0, 1, 2, \dots$, might be considered the definition of F_0 rooted F -cobweb graph Π or in short P if F -sequence has been established. The cobweb poset Π is being thus identified with a **chain of di-bicliques**. The usual convention is to choose $F_0 = 1$. One may relax this constrain, of course. Thus any cobweb one-layer $\langle \Phi_k \rightarrow \Phi_{k+1} \rangle$ is a complete one direction bipartite digraph i.e. a di-biclique. This is how the definition of the F -cobweb graph $\Pi(F)$ or in short P - has emerged.

For infinite cobweb poset Π with the set of vertices $P = V(\Pi)$ one has an obvious $\delta(G) = 2$ domatic vertex partition of this F -cobweb poset.

Definition 8 . A subset D of the vertex set $V(G)$ of a digraph G is called dominating in G , if each vertex of G either is in D , or is adjacent to a vertex of D . Adjacent means that there exists an originating or terminating arc in between the two- any node from G outside D and a node from D .

Definition 9 . A domatic partition of V is a partition of V into dominating sets, and the number of these dominating sets is called the size of such a partition. The domatic number $\delta(G)$ is the maximum size of a domatic partition.

An infinite cobweb poset Π with the set of vertices $P = V(\Pi)$ has the $\delta(\Pi) = 2$ domatic vertex partition, namely a *mod 2* - partition.

$V = V_0 \cup V_1$ where $V_0 = \bigcup_{k=2s+1} \Phi_k$, $s = 0, 1, 2, \dots$ - ("black levels"); and $V_1 = \bigcup_{k=2s} \Phi_k$, $s = 0, 1, 2, \dots$ - ("blue levels").

Note: Natural *mod n* partitions of the cobweb poset's set of vertices $V(\Pi) = P$ (n colours), $P = V_0 \cup V_1 \cup V_2 \cup \dots \cup V_{n-1}$, $V_i = \bigcup_{k=2s+i} \Phi_k$, $s = 0, 1, 2, \dots, i \in \mathbb{Z}_n = 0, 1, \dots, n-1$ for $n > 2$ are **not domatic**.

Cobweb one-layer or more than one-layer subposets $\langle \Phi_k \rightarrow \Phi_{k+m} \rangle \equiv L_{F_k, F_{k+1}} = K_{F_k, F_{k+1}}$, $k = 0, 1, 2, \dots$ have also correspondent, obvious the $\delta(G) = 2$ domatic partitions for $m > 0$.

Combinatorial interpretation.

The crucial and elementary observation now is that an eventual cobweb poset or any combinatorial interpretation of F -binomial coefficients makes sense *not for arbitrary F sequences as F -nomial coefficients should be nonnegative integers* (hybrid sets are not considered here).

Definition 10 . A natural numbers' valued sequence $F = \{n_F\}_{n \geq 0}$, $F_0 = 1$ is called cobweb-admissible iff

$$\binom{n}{k}_F \in \mathbb{N}_0 \quad \text{for} \quad k, n \in \mathbb{N}_0.$$

$F_0 = 0$ being sometimes acceptable as $0_F! \equiv F_0! = 1$.

Incidence coefficients of any reduced incidence algebra of full binomial type [16] immensely important for computer science are computed exactly with their correspondent cobweb-admissible sequences. These include binomial (Newton) or q -binomial (Gauss) coefficients. For other F -nomial coefficients - computed with cobweb admissible sequences - see in what follows after Observation 3.

Problem 1. *Find effective characterizations and/or an algorithm to produce the cobweb admissible sequences i.e. find all examples.*

Right from the definition of P via its Hasse diagram pictures the important observations follow which lead to a specific, new joint combinatorial interpretation of cobweb poset's characteristic binomial-like coefficients.

Observation 1.

The number of maximal chains starting from The Root (level 0_F) to reach any point at the n - th level with n_F vertices is equal to $n_F!$.

Observation 2. ($k > 0$)

The number of all maximal chains in-between $(k + 1)$ - th level Φ_{k+1} and the n - th level Φ_n with n_F vertices is equal to n_F^m , where $m + k = n$.

Indeed. Denote the number of ways to get along maximal chains from *any fixed point* (the leftist for example) in Φ_k to any vertex in $\Phi_n, n > k$ with the symbol

$$[\Phi_k \rightarrow \Phi_n]$$

then obviously we have ($[\Phi_n \rightarrow \Phi_n] \equiv 1$):

$$[\Phi_0 \rightarrow \Phi_n] = n_F!$$

and

$$[\Phi_0 \rightarrow \Phi_k] \times [\Phi_k \rightarrow \Phi_n] = [\Phi_0 \rightarrow \Phi_n].$$

For the purpose of a new joint combinatorial interpretation of F - *sequence - nomial* coefficients (F -nomial - in short) let us consider all finite "max-disjoint" sub-posets rooted at the k - th level at any fixed vertex $\langle r, k \rangle, 1 \leq r \leq k_F$ and ending at corresponding number of vertices at the n - th level ($n = k + m$) where the *max-disjoint* sub-posets are defined below.

Definition 11 . *Two posets are said to be max-disjoint if considered as sets of maximal chains they are disjoint i.e. they have no maximal chain in common. An equipotent copy of P_m ['equip-copy'] is defined as such a maximal chains family equinumerous with P_m set of maximal chains that it constitutes a sub-poset with one minimal element.*

We shall proceed with deliberate notation coincidence anticipating coming observation.

Definition 12 . *Let us denote the number of all mutually max-disjoint equip-copies of P_m rooted at any fixed vertex $\langle j, k \rangle, 1 \leq j \leq k_F$ of k - th level with the symbol*

$$\binom{n}{k}_F.$$

One uses here the customary convention: $\binom{0}{0}_F = 1$ and $\binom{n}{n}_F = 1$.

Compare the above with the Definition 4 and the Definition 10.

The number of ways to reach an upper level from a lower one along any of maximal chains i.e. the number of all maximal chains from the level Φ_{k+1} to the level Φ_n , $k > n$ is equal to

$$[\Phi_k \rightarrow \Phi_n] = n_F^{\frac{m}{F}}.$$

Therefore we have

$$(1) \quad \binom{n}{k}_F \times [\Phi_0 \rightarrow \Phi_m] = [\Phi_k \rightarrow \Phi_n] = n_F^{\frac{m}{F}}$$

where $[\Phi_0 \rightarrow \Phi_m] = m_F!$ counts the number of maximal chains in any equip-copy of P_m . With this in mind we see that the following holds.

Observation 3. $n, k \geq 0$,

Let $n = k + m$. Let F be any cobweb admissible sequence. Then the number of mutually max-disjoint equip-copies i.e. sub-posets equipotent to P_m , rooted at the same **fixed** vertex of k -th level and ending at the n -th level is equal to

$$\begin{aligned} \frac{n_F^{\frac{m}{F}}}{m_F!} &= \binom{n}{m}_F \\ &= \binom{n}{k}_F = \frac{n_F^{\frac{k}{F}}}{k_F!}. \end{aligned}$$

The immediate natural question now is

$$\left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}_{const} = ?$$

i.e. the number of partitions with block sizes all equal to $const = ?$

where here $const = \lambda = m_F!$ and

$$\eta = n_F^{\frac{m}{F}}, \quad \kappa = \binom{n}{k}_F$$

The $const$ indicates that this is the number of set partitions with block sizes all equal to $const$ and we use Knuth notation $\left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}$ for Stirling numbers of the second kind.

From the formula (59) in [4] one infers the Pascal-like matrix answer to the question above.

$$\left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}_{\lambda} = \delta_{\eta, \kappa \lambda} \frac{\eta!}{\kappa! \lambda!^{\kappa}}.$$

This gives us the rough upper bound for the number of tilings (see [1] for Pascal-like triangles) as we arrive now to the following intrinsically related problem.

The partition or tiling Problem 2. Suppose now that F is a cobweb admissible sequence. Let us introduce

$$\sigma P_m = C_m[F; \sigma < F_1, F_2, \dots, F_m >]$$

the equipotent sub-poset obtained from P_m with help of a permutation σ of the sequence $< F_1, F_2, \dots, F_m >$ encoding m layers of P_m thus obtaining the equinumerous sub-poset σP_m with the sequence $\sigma < F_1, F_2, \dots, F_m >$ encoding now m layers of σP_m . Then $P_m = C_m[F; < F_1, F_2, \dots, F_m >]$. Consider the layer $\langle \Phi_k \rightarrow \Phi_n \rangle$, $k < n$, $k, n \in N$ partition into the equal size blocks which are here max-disjoint equi-copies of P_m , $m = n - k + 1$. The question then arises whether and under which conditions the layer may be partitioned with help of max-disjoint blocks of the form σP_m . And how to visualize this phenomenon? It seems to be the question of computer art, too. At first - we already know that an answer to the main question of such tilings existence - for some sequences F - is in affirmative. Whether is it so for all cobweb admissible sequences - we do not know by now. Some computer experiments done by student Maciej Dziemiańczuk [1] are encouraging. More than that. The author of [1] proves tiling's existence for some cobweb-admissible sequences including natural and Fibonacci numbers sequences. He shows also that not all F - designated cobweb posets do admit tiling as defined above. However problems: "how many?" or "find it all" are opened. Let us recapitulate.

Problem 2. Recapitulation. Under which conditions layers may be partitioned with help of max-disjoint blocks of established type σP_m ? Find effective characterizations and/or find an algorithm to produce these partitions.

The problem is intriguing also from the point of view of the art of programming and smart computer experiments. One encounters such situation already looking for simple-minded rough upper bound for the number of tilings (see [1] for Pascal-like triangles). The characteristic feature and effect of calculations and experiments is immediate appearance of Giant Numbers.

GIGANTIC NUMBERS in COBWEB POSET TILING' upper bound PASCAL-LIKE TRIANGLES

Examining in more detail the answer to the already posed question above

$$\left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}_{const} = ?$$

we see that

$$(*) \quad \left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}_{\lambda} = \delta_{\eta, \kappa \lambda} \left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\} = \delta_{\eta, \kappa \lambda} \frac{\eta!}{\kappa!} \sum_{\substack{i_1 + i_2 + \dots + i_{\kappa} = \eta \\ 0 < i_1 = \dots = i_{\kappa} = \lambda}} \frac{1}{i_1! i_2! \dots i_{\kappa}!} = \delta_{\eta, \kappa \lambda} \frac{\eta!}{\kappa! (\lambda!)^{\kappa}}$$

and we constatate the simple-minded recurrence $(**) \equiv (*)$

$$(**) \quad \left\{ \begin{matrix} \eta + \lambda \\ \kappa + 1 \end{matrix} \right\}_{\lambda} = \delta_{\eta + \lambda, (\kappa + 1)\lambda} \frac{(\eta + 1)^{\overline{\lambda}}}{\lambda! (\kappa + 1)} \left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}_{\lambda}; \quad \left\{ \begin{matrix} \lambda \\ 1 \end{matrix} \right\}_{\lambda} = 1$$

resulting in

$$(***) \quad \left\{ \begin{matrix} \kappa \lambda \\ \kappa \end{matrix} \right\}_{\lambda} = \prod_k^{\kappa} \Phi_{\lambda}(k) \equiv [\Phi_{\lambda}(\kappa)]!$$

which we shall call Φ_λ - factorial, where

$$\Phi_\lambda(\kappa) = \frac{(\kappa\lambda)^\lambda}{\lambda!\kappa}$$

Résumé. The number of κ - block partitions of $\kappa\lambda$ - numerous ensemble with block sizes all equal to $\lambda = m_F!$ is equal to its Φ_λ - factorial.

Experimental comparison. Hereby we show and compare few upper bound numbers $\Phi_\lambda(\kappa)$ with the number $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}_F$ of **all** different tilings of the layer $\langle \Phi_k \rightarrow \Phi_n \rangle$ with σP_m blocks as computer experiments show [1]. Except for obvious cases $\Phi_\lambda(\kappa) = 1$ and very few first rows of upper-bound F - triangles (zeros being not displayed), we are facing - face to face - enormously Gigantic Numbers as illustrated below via quotations from [1].

$$\begin{array}{c} \text{Kwasniewski Cobweb triangle} \quad \Phi_\lambda(\kappa)! = \left\{ \begin{smallmatrix} \eta \\ \kappa \end{smallmatrix} \right\}_\lambda \\ F - \text{Natural numbers} \\ 1 \\ 1 \quad 1 \\ 1 \quad 15 \quad 1 \\ 1 \quad \frac{24!}{4! \cdot 6!^4} \quad 10395 \quad 1 \\ 1 \quad \frac{120!}{5! \cdot 24!^{15}} \quad \frac{60!}{10! \cdot 6!^{10}} \quad \frac{20!}{10! \cdot 2!^{10}} \quad 1 \\ 1 \quad \frac{720!}{6! \cdot 120!^6} \quad \frac{360!}{15! \cdot 24!^{15}} \quad \frac{120!}{20! \cdot 6!^{20}} \quad \frac{30!}{15! \cdot 2!^{15}} \quad 1 \\ 1 \quad \frac{5040!}{7! \cdot 720!^7} \quad \frac{2520!}{21! \cdot 120!^{21}} \quad \frac{840!}{35! \cdot 24!^{35}} \quad \frac{210!}{35! \cdot 6!^{35}} \quad \frac{42!}{21! \cdot 2!^{21}} \quad 1 \\ \dots \end{array}$$

Fig.7. Display of Natural numbers' case - calculated upper bound.

$$\begin{array}{c} \text{computer experiments} \\ \text{Cobweb poset tiling triangle for} \quad \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}_F \\ F - \text{Natural numbers} \\ 1 \\ 1 \quad 1 \\ 1 \quad 4 \quad 1 \\ 1 \quad 32 \quad 132 \quad 1 \\ 1 \quad 386 \quad 411168 \quad 44928 \quad 1 \\ 1 \quad 12142 \quad \quad \quad 243319680 \quad 1 \\ \dots \end{array}$$

Fig.8. Display of Natural numbers case - number of tilings from experiment.

$$\begin{array}{c}
\text{Kwasniewski Cobweb triangle } \Phi_{\lambda}(\kappa) = \left\{ \begin{matrix} \eta \\ \kappa \end{matrix} \right\}_{\lambda} \\
F - \text{Fibonacci numbers} \\
\begin{array}{ccccccc}
& & & 1 & & & \\
& & 1 & & 1 & & \\
& 1 & & 1 & & 1 & \\
& & 1 & & 15 & & 1 \\
& 1 & & \frac{30!}{5! \cdot 6!^5} & & \frac{30!}{15! \cdot 2!^{15}} & 1 \\
& & 1 & & \frac{240!}{8! \cdot 30!^8} & & \frac{240!}{40! \cdot 6!^{40}} & & \frac{120!}{60! \cdot 2!^{60}} & 1 & 1 \\
1 & & \frac{3120!}{13! \cdot 240!^{13}} & & \frac{3120!}{104! \cdot 30!^{104}} & & \frac{1560!}{260! \cdot 6!^{260}} & & \frac{520!}{260! \cdot 2!^{260}} & 1 & 1
\end{array}
\end{array}$$

...

Fig.9. Display of Fibonacci numbers' case - calculated upper bound.

$$\begin{array}{c}
\text{computer experiments} \\
\text{Cobweb poset tiling triangle for } \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_F \\
F - \text{Fibonacci numbers} \\
\begin{array}{ccccccc}
& & & 1 & & & \\
& & 1 & & 1 & & \\
& 1 & & 1 & & 1 & \\
& & 1 & & 4 & & 1 \\
1 & & 136 & & 7176824 & & 1
\end{array}
\end{array}$$

...

Fig.10. Display of Natural numbers case - number of tilings from experiment.

Coming over to the last problem announced let us note that the Observation 3. provides us with the *new* combinatorial interpretation of the immense class of all classical F – *nomial* coefficients including binomial or Gauss q -binomial ones or Konvalina generalized binomial coefficients of the first and of the second kind [4] - which include Stirling numbers too. All these F -nomial coefficients naturally are computed with their correspondent cobweb-admissible sequences. More than that - the vast ‘umbral’ family of F -sequences [10,11,12,13,6] includes also those which shall be called “*GCD-morphic*” sequences. This means that $GCD[F_n, F_m] = F_{GCD[n,m]}$ where GCD stays for Greatest Common Divisor.

Definition 13 . *The sequence of integers $F = \{n_F\}_{n \geq 0}$ is called the *GCD-morphic sequence* if $GCD[F_n, F_m] = F_{GCD[n,m]}$ where GCD stays for Greatest Common Divisor operator.*

The Fibonacci sequence is a much nontrivial [11,12,6] guiding example of GCD-morphic sequence. Of course *not all* incidence coefficients of reduced incidence

algebra of full binomial type are computed with GCD-morphic sequences however these or that - if computed with the cobweb correspondent admissible sequences all are given the new, joint cobweb poset combinatorial interpretation via Observation 3. More than that - in [8] a prefab-like combinatorial description of cobweb posets is being served with corresponding generalization of the fundamental exponential formula.

Question: which of these above mentioned sequences are GCD-morphic sequences?

GCD-morphism Problem. Problem III. *Find effective characterizations and/or an algorithm to produce the GCD-morphic sequences i.e. find all examples.*

The recent papers on DAGs related to this article and its clue references [8,9] apart from [1] are [7] and [14].

For the by now remark. For resented results on cobweb posets -up to the January 2008 - see [19]. The *poli – ary* relations' (as in relational database models) point of view on cobweb posets and their KoDAGs representatives see [20] from 21 of December 2008 and for thus resulting natural open problems see then [21].

Acknowledgements The author appraises much Maciej Dziemiańczuk's computer experiments founded on his vivid-active and effective interest in cobweb posets' now our joint investigations. I am also indebted for his successful indication of an important misprint. The author also appraises Maciej Dziemiańczuk now Gdańsk University Student's *TeX – nology* aid.

The author expresses **solemnly** his gratitude to Dr Ewa Krot-Sieniawska for her several years' cooperation until **she was kicked out by Białystok University authorities exactly on the day she had defended her Rota and KoDAs related dissertation with distinction**. Innocent was thus penalized **because of me** unmasking in public misconducts, infringements and contraventions.

References

- [1] M. Dziemiańczuk, *On cobweb posets tiling problem*, Adv. Stud. Contemp. Math. volume 16 (2), 2008 (April) pp. 219-233, *arXiv* : *math.Co/0709.4263*, 4 Oct 2007.
- [2] Curtis Greene, *www.haverford.edu/math/cgreene/posets/posetgallery.html*.
- [3] Hasse Diagram, <http://mathworld.wolfram.com/HasseDiagram.html>.
- [4] Charles Jordan *On Stirling Numbers* Thoku Math. J. **37** (1933), 254-278.
- [5] J. Konvalina , *A Unified Interpretation of the Binomial Coefficients, the Stirling Numbers and the Gaussian Coefficients* The American Mathematical Monthly **107** (2000), 901-910.
- [6] E. Krot, *An Introduction to Finite Fibonomial Calculus*, CEJM 2(5) (2005) 754-766.
- [7] E. Krot, *The first ascent into the Fibonacci Cob-web Poset*, Adv. Stud. Contemp. Math. **11** (2) (2007) 179-184.

- [8] A. K. Kwaśniewski, *Cobweb posets as noncommutative prefabs* Adv. Stud. Contemp. Math. **14** (1) (2007) 37-47.
- [9] A. K. Kwaśniewski: *First Observations on Prefab Posets' Whitney Numbers* Advances in Applied Clifford Algebras **Vol.18**, Number 1 / February, **2008** 57-73. arXiv: math.CO /0802.1696 cs.DM
- [10] A. K. Kwaśniewski *Main theorems of extended finite operator calculus* Integral Transforms and Special Functions, **14** (6) (2003) 499-516.
- [11] A. K. Kwaśniewski, *The logarithmic Fib-binomial formula*, Advanced Stud. Contemp. Math. **9** No 1 (2004) 19-26.
- [12] A. K. Kwaśniewski, *Fibonomial cumulative connection constants*, Bulletin of the ICA **44** (2005) 81-92.
- [13] A. K. Kwaśniewski, *On umbral extensions of Stirling numbers and Dobinski-like formulas* Advanced Stud. Contemp. Math. **12**(2006) no. 1, pp.73-100.
- [14] Anatoly D. Plotnikov, *About presentation of a digraph by dim 2 poset*, Adv. Stud. Contemp. Math. **12** (1) (2006) 55-60
- [15] Bruce E. SAGAN *Mobius Functions of Posets* (Lisbon lectures)IV: *Why the Characteristic Polynomial factors* June 28 **2007** <http://www.math.msu.edu/>
- [16] E. Spiegel, Ch. J. O'Donnell *Incidence algebras* Marcel Dekker, Inc., Basel, 1997.
- [17] Richard P. STANLEY, *Hyperplane Arrangements*, Proc. Nat. Acad. Sci. **93** (1996), 2620-2625. *An Introduction to Hyperplane Arrangements* www.math.umn.edu/~ezra/PCMI2004/stanley.pdf
- [18] M. Ward: *A calculus of sequences*, Amer.J.Math. **58** (1936) 255-266.
- [19] A. Krzysztof Kwaśniewski, M. Dziemiańczuk, *Cobweb posets - Recent Results*, Adv. Stud. Contemp. Math. vol. 16 (2) April 2008 . pp. 197-218 ; arXiv: math. /0801.3985 Fri, 25 Jan 2008 17:01:28 GMT
- [20] A. Krzysztof Kwaśniewski, *Cobweb Posets and KoDAG Digraphs are Representing Natural Join of Relations, their diBigraphs and the Corresponding Adjacency Matrices*, arXiv:math/0812.4066v1 ,[v1] Sun, 21 Dec 2008 23:04:48 GMT
- [21] A. Krzysztof Kwaśniewski, *Some Cobweb Posets Digraphs' Elementary Properties and Questions*, arXiv:0812.4319v1 ,[v1] Tue, 23 Dec 2008 00:40:41 GMT